arXiv:1407.7603v1 [math.PR] 28 Jul 2014

ON SOME SMOOTHENING EFFECTS OF THE TRANSITION SEMIGROUP OF A LÉVY PROCESS

ZHAO DONG, SZYMON PESZAT, AND LIHU XU

ABSTRACT. Let (P_t) be the transition semigroup of a Lévy process L taking values in a Hilbert space H. Let ν be the Lévy measure of L. It is shown that for any bounded and measurable function f,

$$\int_H |P_t f(x+y) - P_t f(x)|^2 \, \nu(\mathrm{d}y) \leq \frac{1}{t} P_t f^2(x) \qquad \text{for all } t > 0,\ x \in H.$$

As ν can be infinite this formula establishes some smoothening effect of the semigroup (P_t). In the paper some applications of the formula will be presented as well.

1. INTRODUCTION

Let (X_t) be the solution to an SDE on a Hilbert space H driven by a non-degenerate Wiener process W. Let

$$P_t f(x) = \mathbb{E}\left(f(X_t)|X_0 = x\right), \qquad f \in B_b(H),\ t \geq 0,$$

be the corresponding transition semigroup defined on the space of bounded measurable functions $B_b(H)$. Then the following Bismut–Elworthy–Li formula holds (see [4] or [8])

$$\langle \nabla P_t f(x), h\rangle_H = \frac{1}{t}\mathbb{E}\left(f(X_t)\int_0^t K(s;h)\mathrm{d}W_s|X_0 = x\right),$$

where $K(s,h)$ is an adapted stochastic processes independent of f. This formula implies the strong Feller property of (P_t), and therefore is very useful for studying its ergodic properties. For other applications we refers readers to e.g. [14, 2, 5].

In this paper, we prove that the transition semigroup (P_t) of a Lévy process L taking values in a Hilbert space H exhibits some smoothening effects, namely P_t transforms $B_b(H)$ into the intersection of domains of some non-local operators (see Theorem 3.1 for more details). The proof of Theorem 3.1 is very simple and follows [4]. Next, see Corollary 3.2, we will show that

$$\int_H |P_t f(x+y) - P_t f(x)|^2 \, \nu(\mathrm{d}y) \leq \frac{1}{t} P_t f^2(x) \quad \text{for } f \in B_b(H),\ t > 0,\ x \in H. \tag{1.1}$$

Above ν is the Lévy measure of L. Note that if ν is infinite, then for any open ball $B_\varepsilon(0)$ with the center at 0 and radius $\varepsilon > 0$ one has $\nu\left(B_\varepsilon(0)\right) = +\infty$. Therefore (1.1) means that for any $x \in H$, $P_t f(x+y)$, $y \in B_\varepsilon(0)$, is in a certain sense close to $P_t f(x)$.

As applications of our general result, we obtain a short time estimate for the semigroup generated by fractional Laplacian and a 'fractional gradient' estimate for the perturbed

ZD is supported by Key Laboratory of Random Complex Structures and Data Science, Academy of Mathematics and Systems Science, CAS (No.2008DP173182), NSFC(No. 11271356), NSFC(No. 10721101), 973 Program (2011CB808000). LX is supported by the grant SRG2013-00064-FST. SP is supported by Polish National Science Center grant DEC2013/09/B/ST1/03658. The authors would like to gratefully thank Prof. Zhen-Qing Chen and Dr. Jieming Wang for stimulating discussions, and Profs. Jiahong Wu and Zhifei Zhang for their helpful discussions on fractional Laplacian and Campanato's theorem.

stable type stochastic systems considered in [3]. Finally using generalized Campanato's theorem, we calculate modulus of continuity of the transition semigroups of 'log α-stable' processes.

The paper is organized as follows: the next section includes some preliminary facts on Lévy processes. The main general results are formulated in Section 3. The last three sections are devoted to applications. In the appendix we prove the generalized Campanato theorem of harmonic analysis.

2. Preliminary facts on Lévy processes

We shall recall here some preliminary facts on Lévy processes (for details see e.g. [1, 9, 10]). Let $(L_t)_{t\geq 0}$ be an H-valued Lévy process. It is well known that there is a vector $m \in H$, a symmetric positive definite trace class operator $Q\colon H \mapsto H$, and a Borel measure ν on H satisfying

$$\nu(\{0\}) = 0, \qquad \int_H 1 \wedge |y|_H^2 \nu(\mathrm{d}y) < +\infty, \tag{2.1}$$

such that

$$\mathbb{E}\, \mathrm{e}^{\mathrm{i}\langle x, L_t\rangle_H} = \mathrm{e}^{-t\psi(x)}, \qquad x \in H,$$

where the so-called *Lévy symbol* ψ of (L_t) is given by

$$\psi(x) = i\langle x, m\rangle_H + \frac{1}{2}\langle Qx, x\rangle_H + \int_H \left[\mathrm{e}^{\mathrm{i}\langle x,y\rangle_H} - 1 - \mathrm{i}\langle x, y\rangle_H \mathbf{1}_{\{|y|_H\leq 1\}}\right] \nu(\mathrm{d}y).$$

We call ν the *Lévy measure* of L and (m, Q, ν) the *generating triplet* of L.

The *Poisson random measure* associated with (L_t) is defined by

$$N(t, \Gamma) := \sum_{s\in(0,t]} \mathbf{1}_\Gamma(L_s - L_{s-}), \qquad \Gamma \in \mathcal{B}(H),\ t > 0,$$

and the *compensated Poisson random measure* is given by

$$\tilde{N}(t, \Gamma) = N(t, \Gamma) - t\nu(\Gamma).$$

By the *Lévy–Khinchin decomposition* (cf. [1, p.108, Theorem 2.4.16] or [9, p. 53, Theorem 4.23]), one has

$$L_t = mt + W_Q(t) + \int_{\{0<|x|_H\leq 1\}} x\tilde{N}(\mathrm{d}t, \mathrm{d}x) + \int_{\{|x|_H>1\}} xN(\mathrm{d}t, \mathrm{d}x), \qquad t \geq 0,$$

where W_Q is a Wiener process in H with covariance operator Q.

Let $(\mathfrak{F}_t)$ be the filtration generated by (L_t), and let us denote by $\mathcal{L}^2_{loc}$ the space of all predictable stochastic process ψ satisfying

$$\mathbb{E}\int_0^t \int_H |\psi(s, y)|_H^2 \nu(\mathrm{d}y)\mathrm{d}s < \infty \qquad \text{for } t > 0.$$

Then for any $\psi \in \mathcal{L}^2_{loc}$ the stochastic integral $\int_0^t \int_H \psi(s, y)\tilde{N}(\mathrm{d}s, \mathrm{d}y)$ is a well-defined square integrable and mean zero martingale. Moreover, the following Itô isometry holds (see e.g. [1, p. 200] or [9, Section 8.7])

$$\begin{aligned} &\mathbb{E}\left[\int_0^t \int_H \psi(s, y)\tilde{N}(\mathrm{d}s, \mathrm{d}y) \int_0^t \int_H \varphi(s, y)\tilde{N}(\mathrm{d}s, \mathrm{d}y)\right] \\ &\quad = \mathbb{E}\int_0^t \int_H \psi(s, y)\varphi(s, y)\nu(\mathrm{d}y)\mathrm{d}s, \qquad \psi, \varphi \in \mathcal{L}^2_{loc}. \end{aligned} \tag{2.2}$$

Let $L = (L_t)$ be a Lévy process with a generating triplet (m, Q, ν). Consider the Markov family

$$L_t^x = x + L_t, \qquad t \geq 0,\ x \in H. \tag{2.3}$$

Its transition semigroup (P_t) is given as follows

$$P_t f(x) = \mathbb{E}\, f(L_t^x), \qquad t \geq 0,\ x \in H,\ f \in B_b(H). \tag{2.4}$$

Observe that (P_t) is C_0 on the space $UC_b(H)$ of uniformly continuous bounded functions on H, see [9, p. 80]. Moreover, the domain of its generator $\mathcal{L}$ contains the space $UC_b^2(H)$, and

$$\mathcal{L}f(x) = \langle Df(x), m\rangle_H + \frac{1}{2}\operatorname{Trace} QD^2 f(x) + \int_H \left(f(x+y) - f(x) - \mathbf{1}_{\{|x|_H<1\}}\langle Df(x), y\rangle_H\right)\nu(\mathrm{d}y), \qquad f \in UC_b^2(H),\ t \geq 0,\ x \in H.$$

3. Main results

Let L be a Lévy process on H with the generation triplet (m, Q, ν). Let (L_t^x), $t \geq 0$, $x \in H$ be the Markov family given by (2.3), and let (P_t) given by (2.4) be the transition semigroup of (L_t^x).

Given $q \in L^2(H, \mathcal{B}(H), \nu)$, define

$$\mathcal{D}_q := \left\{ f \in B_b(H) \colon \sup_{x\in H} \int_H |f(x+y) - f(x)|\,|q(y)|\nu(\mathrm{d}y) < \infty \right\}.$$

Next, let

$$A_q f(x) := \int_H [f(x+y) - f(x)]\, q(y)\nu(\mathrm{d}y) \qquad \text{for } f \in \mathcal{D}_q,\ x \in H.$$

Taking into account (2.1), we see that $C_b^1(H) \subset \mathcal{D}_q$, A_q is a bounded linear operator form $C_b^1(H)$ into $B_b(H)$ and

$$\begin{aligned}\|A_q f\|_\infty &:= \sup_{x\in H} |A_q f(x)| \\ &\leq 2\|f\|_\infty \left(\int_{|y|_H\geq 1} \nu(\mathrm{d}y)\right)^{\frac{1}{2}} \left(\int_{|y|_H\geq 1} q^2(y)\nu(\mathrm{d}y)\right)^{\frac{1}{2}} \\ &\quad + \|Df\|_\infty \left(\int_{\{|y|_H<1\}} |y|_H^2 \nu(\mathrm{d}y)\right)^{\frac{1}{2}} \left(\int_{\{|y|_H<1\}} q^2(y)\nu(\mathrm{d}y)\right)^{\frac{1}{2}} < \infty.\end{aligned}$$

Theorem 3.1. *Let $q \in L^2(H, \mathcal{B}(H), \nu)$. Then for all $t > 0$ and $f \in B_b(H)$, $P_t f \in \mathcal{D}_q$ and*

$$|A_q P_t f(x)|^2 \leq \frac{1}{t} P_t f^2(x) \int_H q^2(y)\nu(\mathrm{d}y) \qquad \textit{for all } x \in H.$$

Proof. Assume that $f \in UC_b^2(H)$. Applying Itô's formula (see e.g. [1] or [9]) we obtain

$$\begin{aligned} f(L_t^x) - P_t f(x) = &- \int_0^t \mathcal{L} P_{t-s} f(L_s^x) \mathrm{d}s + \int_0^t P_{t-s} \mathcal{L} f(L_s^x) \mathrm{d}s \\ &+ \int_0^t \int_H \left[P_{t-s} f(L_s^x + y) - P_{t-s} f(L_s^x) \right] \tilde{N}(\mathrm{d}y, \mathrm{d}s) \\ &+ \int_0^t \langle D P_{t-s} f(L_s^x), \mathrm{d}W_Q(s) \rangle_H \\ = &\int_0^t \int_H \left[P_{t-s} f(X_s + y) - P_{t-s} f(X_s) \right] \tilde{N}(\mathrm{d}y, \mathrm{d}s) \\ &+ \int_0^t \langle D P_{t-s} f(L_s^x), \mathrm{d}W_Q(s) \rangle_H, \end{aligned} \tag{3.1}$$

where the last equality is because $\mathcal{L} P_t f = P_t \mathcal{L} f$ for $f \in UC_b^2(H)$. Multiplying the both sides of (3.1) by

$$\int_0^t \int_H q(y) \tilde{N}(\mathrm{d}y, \mathrm{d}s)$$

and taking into account (2.2), we further get

$$\begin{aligned} \mathbb{E}\left[f(L_t^x) \int_0^t \int_H q(y) \tilde{N}(\mathrm{d}y, \mathrm{d}s) \right] &= \mathbb{E} \int_0^t \int_H \left[P_{t-s} f(L_s^x + y) - P_{t-s} f(L_s^x) \right] q(y) \nu(\mathrm{d}y) \mathrm{d}s \\ &= \int_0^t \int_H \left[P_s P_{t-s} f(x + y) - P_s P_{t-s} f(x) \right] q(y) \nu(\mathrm{d}y) \mathrm{d}s \\ &= t \int_H \left[P_t f(x + y) - P_t f(x) \right] q(y) \nu(\mathrm{d}y) = t A_q P_t f(x). \end{aligned}$$

Thus, by the Hölder inequality and Itô isometry we obtain

$$\begin{aligned} t \left| A_q P_t f(x) \right| &\le \left(\mathbb{E}\, f^2(L_t^x) \right)^{1/2} \left(\int_0^t \int_H q^2(y) \nu(\mathrm{d}y) \mathrm{d}s \right)^{1/2} \\ &\le \left(P_t f^2(x) \right)^{1/2} t^{1/2} \left(\int_H q^2(y) \nu(\mathrm{d}y) \right)^{1/2}. \end{aligned}$$

Thus the desired estimate holds for any $f \in UC_b^2(H)$. Assume that $f \in B_b(H)$. Let $x \in H$. Then there is a sequence $(f_n) \subset UC_b^2(H)$ such that

$$\lim_{n \to \infty} P_t f_n^2(x) = P_t f^2(x),$$

and

$$\lim_{n \to \infty} P_t f_n(x + y) = P_t f(x + y) \quad \text{for } \nu \text{ almost all } y.$$

Consequently, the desired estimate for f follows from the Fatou lemma. □

Corollary 3.2. *For arbitrary $f \in B_b(H)$ we have*

$$\int_H \left| P_t f(x + y) - P_t f(x) \right|^2 \nu(\mathrm{d}y) \le \frac{1}{t} P_t f^2(x), \qquad x \in H,\ t > 0.$$

Proof. Since

$$\int_H \left| P_t f(x + y) - P_t f(x) \right|^2 \nu(\mathrm{d}y) = \sup \left\{ \left| A_q P_t f(x) \right|^2, \quad q \colon \int_H q^2(y) \nu(\mathrm{d}u) \le 1 \right\}$$

the estimate follows from Theorem 3.1. □

Given $f \in B_b(H)$ we define the difference operator $\nabla^n_{y_1,\dots,y_n} f(x)$, $x, y_1, \dots, y_n \in H$ putting

$$\nabla_y f(x) = f(x+y) - f(x),$$
$$\nabla^{n+1}_{y_1,\dots,y_{n+1}} f(x) = \nabla_{y_{n+1}} \left(\nabla^n_{y_1,\dots,y_n} f\right)(x).$$

Corollary 3.3. *For any $f \in B_b(H)$ and $n \in \mathbb{N}$,*

$$\sup_{x\in H} \int_H \dots \int_H \left|\nabla_{y_1,\dots,y_n} (P_t f)(x)\right|^2 \nu(\mathrm{d}y_1) \dots \nu(\mathrm{d}y_n) \le \left(\frac{n}{t}\right)^n \|f\|^2_\infty.$$

Proof. It is enough to show the estimate for $f \in UC^2_b(H)$. Let $q_1, \dots, q_n \in L^2(H, \mathcal{B}(H), \nu)$. We claim that the operators A_q and P_s commute. Indeed, by the Feynman–Kac representation of P_t, the Fubini theorem and the fact $X_t(x) + y = X_t(x+y)$, for all $f \in UC^2_b(\mathbb{R}^d)$ we have

$$\begin{aligned} P_t A_q f(x) &= \mathbb{E}\, A_q f(X_t(x)) = \mathbb{E} \int_H \left(f(X_t(x)+y) - f(X_t(x))\right) q(y)\nu(\mathrm{d}y) \\ &= \int_H \left[\mathbb{E} f(X_t(x)+y) - \mathbb{E} f(X_t(x))\right] q(y)\nu(\mathrm{d}y) \\ &= \int_H \left[P_t f(x+y) - P_t f(x)\right] q(y)\nu(\mathrm{d}y) \\ &= A_q P_t f(x). \end{aligned}$$

Thus by Theorem 3.1,

$$\begin{aligned} \|A_{q_1} \dots A_{q_n} P_t f\|^2_\infty &= \sup_{x\in H} |A_{q_1} \dots A_{q_n} P_t f(x)|^2 \\ &= \sup_{x\in H} \left|(A_{q_1} P_{t/n}) \dots (A_{q_n} P_{t/n}) f(x)\right|^2 \\ &\le \frac{n}{t} \|(A_{q_1} P_{t/n}) \dots (A_{q_n} P_{t/n}) f\|^2_\infty. \end{aligned}$$

□

4. Application 1: Short time behaviour of the semigroup

We shall study the fractional gradient estimate of α-stable and truncated α-stable processes. In this and next sections $H = \mathbb{R}^d$. The norm on $\mathbb{R}^d$ will be denoted by $|\cdot|$.

Theorem 4.1. *Let the Lévy measure ν be of the form*

$$\nu(\mathrm{d}x) = \frac{1}{|x|^{d+\alpha}} \mathbf{1}_{\{|x|<K\}} \mathrm{d}x,$$

where $\alpha \in (0,2]$ and $K \in (0,\infty]$. Then for any $\beta \in (\alpha/2, \alpha)$ we have

$$\|(-\Delta)^{\frac{\alpha-\beta}{2}} P_t f\|_\infty \le C(1 + t^{-1/2})\|f\|_\infty, \qquad \forall\, t > 0,\ f \in B_b(\mathbb{R}^d), \tag{4.1}$$

where $C = C_{\alpha,\beta}$ only depends on α and β.

Proof. Without any loss of generality, we may assume that $K = 1$. Choose q such that

$$q(y) = |y|^\beta \qquad \forall\, |y| \le 1, \qquad \int_{\{|y|>1\}} q^2(y)\nu(\mathrm{d}y) < \infty.$$

It is easy to see that

$$\int_{\{|y|\le 1\}} q^2(y)\nu(\mathrm{d}y) = \int_{\{|y|\le 1\}} |y|^{-d-\alpha+2\beta} \mathrm{d}y \le C_{\alpha,\beta}, \tag{4.2}$$

where $C_{\alpha,\beta} > 0$ depends on α, β. Therefore, $q \in L^2(\nu) := L^2(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d), \nu)$.

Observe that

$$A_q f(x) = \int_{\{|y|\le 1\}} \frac{f(x+y)-f(x)}{|y|^{\alpha+d-\beta}}\mathrm{d}y + \int_{\{|y|>1\}} [f(x+y)-f(x)]\, q(y)\nu(\mathrm{d}y)$$

and

$$\begin{aligned}\left|(-\Delta)^{\frac{\alpha-\beta}{2}} f(x)\right| &= \left|\int_{\mathbb{R}^d} \frac{f(x+y)-f(x)}{|y|^{\alpha+d-\beta}}\mathrm{d}y\right| \\ &\le \left|\int_{\{|y|>1\}} \frac{f(x+y)-f(x)}{|y|^{\alpha+d-\beta}}\mathrm{d}y\right| + \left|\int_{\{|y|\le 1\}} \frac{f(x+y)-f(x)}{|y|^{\alpha+d-\beta}}\mathrm{d}y\right|.\end{aligned}$$

It is easy to see that

$$\left|\int_{\{|y|>1\}} \frac{f(x+y)-f(x)}{|y|^{\alpha+d-\beta}}\mathrm{d}y\right| \le c_{\alpha,\beta}\|f\|_\infty,$$

where $c_{\alpha,\beta} > 0$ depends on α, β, and that

$$\begin{aligned}\left|\int_{\{|y|\le 1\}} \frac{f(x+y)-f(x)}{|y|^{\alpha+d-\beta}}\mathrm{d}y\right| &\le |A_q f(x)| + \left|\int_{\{|y|>1\}} [f(x+y)-f(x)]\, q(y)\nu(\mathrm{d}y)\right| \\ &\le t^{-1/2}\|f\|_\infty \|q\|_{L^2(\nu)} + 2\|f\|_\infty \|q\|_{L^2(\nu)},\end{aligned}$$

where the last inequality follows from Theorem 3.1 and Hölder's inequality. Collecting the previous inequalities, we get the desired one. □

5. Application 2: Estimate for a perturbed dynamics studied in [3]

Let $X_t(x)$ be the value at t of the solution to the following stochastic differential equation

$$\mathrm{d}X_t = b(X_{t-})\mathrm{d}Y_t + \mathrm{d}Z_t, \qquad X_0 = x, \tag{5.1}$$

where $b \in C_b^2(\mathbb{R}^d)$ and Z_t, Y_t are both symmetric stable processes with the parameters $\alpha, \beta \in (0,2)$. Our result below is also true if b is a bounded measurable function, to avoid the complicated differentiability issue and stress the idea, we assume $b \in UC_b^2(\mathbb{R}^d)$.

Eq. (5.1) in more general setting has been studied by Chen and Wang ([3]) and has a unique weak solution. Let

$$P_t f(x) = \mathbb{E} f(X_t(x)), \qquad \forall\, f \in B_b(\mathbb{R}^d),$$

be the corresponding transition semigroup. The semigroup is C_0 on the space $UC_b(\mathbb{R}^d)$. Let $\mathcal{L}$ be the generator of (P_t) considered on $UC_b(\mathbb{R}^d)$. It is well known that $UC_b^2(\mathbb{R}^d) \subset \mathrm{Dom}\,(\mathcal{L})$, and that for all $f \in UC_b^2(\mathbb{R}^d)$,

$$\mathcal{L} f = -(-\Delta)^{\alpha/2} f - |b(x)|^\beta (-\Delta)^{\beta/2} f,$$

and the following backward Kolmogorov equation holds

$$\partial_t P_t f = \mathcal{L} P_t f. \tag{5.2}$$

We shall use (4.1) to show some properties of the associated backward Kolmogorov equation.

Theorem 5.1. *If $\beta \in (0, \alpha/2)$, then there exists a $t_0 \in (0,1)$ depending on α, β and $\|b\|_\infty$, such that for any $f \in UC_b(\mathbb{R}^d)$,*

$$\begin{aligned}&\|(-\Delta)^{\beta/2} P_t f\|_\infty \le C_1 t^{-1/2}\|f\|_\infty, \qquad t \le t_0, \\ &\|(-\Delta)^{\beta/2} P_t f\|_\infty \le C_2 \|f\|_\infty, \qquad t > t_0,\end{aligned} \tag{5.3}$$

where C_1, C_2 depend on α, β and t_0.

Proof. Since $UC_b^2(\mathbb{R}^d)$ is dense in $UC_b(\mathbb{R}^d)$ and $(-\Delta)^{\beta/2}$ is closable, it suffices to show (5.3) for $f \in UC_b^2(\mathbb{R}^d)$.

For any $f \in UC_b^2(\mathbb{R}^d)$, define $P_t^0 f(x) = \mathbb{E}[f(Z_t + x)]$. It satisfies

$$\partial_t P_t^0 f = -(-\Delta)^{\alpha/2} P_t^0 f. \tag{5.4}$$

$(P_t^0)_{t\geq 0}$ can be extended to a Markov C_0-semigroup on $UC_b(\mathbb{R}^d)$. Thanks to (5.2) and (5.4), using the classical Duhamel principle we obtain

$$P_t f(x) = P_t^0 f(x) - \int_0^t P_{t-s}^0 [|b|^\beta (-\Delta)^{\beta/2} P_s f](x) \mathrm{d}s. \tag{5.5}$$

Since $b \in UC_b^2(\mathbb{R}^d)$ and $f \in UC_b^2(\mathbb{R}^d)$, $P_t f \in UC_b^2(\mathbb{R}^d)$ and $P_t^0 f \in UC_b^2(\mathbb{R}^d)$ both hold. By (4.1), we have

$$\|(-\Delta)^{\beta/2} P_t^0 f\|_\infty \leq C(1 + t^{-1/2}) \|f\|_\infty \leq C t^{-1/2} \|f\|_\infty, \qquad \forall\, t < 1, \tag{5.6}$$

which, together with (5.5), yields

$$\begin{aligned}
\|(-\Delta)^{\beta/2} P_t f\|_\infty &\leq C t^{-1/2} \|f\|_\infty + C \int_0^t (t-s)^{-1/2} \|b\|_\infty^\beta \|(-\Delta)^{\beta/2} P_s f\|_\infty \mathrm{d}s \\
&\leq C t^{-1/2} \|f\|_\infty + C \int_0^t (t-s)^{-1/2} \|b\|_\infty^\beta \|(-\Delta)^{\beta/2} P_s f\|_\infty \mathrm{d}s \\
&= C t^{-1/2} \|f\|_\infty + C \int_0^t (t-s)^{-1/2} s^{-1/2} \|b\|_\infty^\beta s^{1/2} \|(-\Delta)^{\beta/2} P_s f\|_\infty \mathrm{d}s.
\end{aligned}$$

Define

$$L_T := \sup_{0\leq t\leq T} t^{1/2} \|(-\Delta)^{\beta/2} P_t f\|_\infty$$

with $T > 0$ to be chosen later. From the previous inequality we have

$$\begin{aligned}
L_T &\leq C\|f\|_\infty + C T^{\frac{1}{2}} \sup_{0\leq t\leq T} \int_0^t s^{-\frac{1}{2}} (t-s)^{-\frac{1}{2}} \mathrm{d}s L_T \\
&= C\|f\|_\infty + C B(3/2, 3/2) T^{\frac{1}{2}} \|b\|_\infty^\beta L_T,
\end{aligned} \tag{5.7}$$

where B is the beta function. Choose a $t_0 \in (0,1)$ (depending on α, β, and $\|b\|_\infty^\beta$) such that $C B(3/2,3/2) t_0^{\frac{1}{2}} \|b\|_\infty^\beta < \frac{1}{2}$, we obtain

$$L_{t_0} \leq 2C\|f\|_\infty.$$

This immediately implies the first estimate in the theorem.

For the second estimate, taking $P_{t_0/2} f$ rather than f as the initial data, by the same procedure as above we have

$$\begin{aligned}
\|(-\Delta)^{\beta/2} P_t f\|_\infty &\leq C_1 \left(t - \frac{t_0}{2}\right)^{-\frac{1}{2}} \|P_{t_0/2} f\|_\infty \\
&\leq C_1 \left(t - \frac{t_0}{2}\right)^{-\frac{1}{2}} \|f\|_\infty, \qquad \forall\, t \in (t_0/2, 3t_0/2).
\end{aligned}$$

Therefore

$$\|(-\Delta)^{\beta/2} P_t f\|_\infty \leq C_1 \left(\frac{t_0}{2}\right)^{-\frac{1}{2}} \|f\|_\infty, \qquad \forall\, t \in (t_0, 3t_0/2).$$

Now taking $P_{t_0}f$ as the initial data, we obtain

$$\|(-\Delta)^{\beta/2}P_tf\|_\infty \le C_1\left(\frac{t_0}{2}\right)^{-\frac{1}{2}}\|f\|_\infty, \qquad \forall\, t\in(3t_0/2, 2t_0).$$

Iterating the above argument, we finally get

$$\|(-\Delta)^{\beta/2}P_tf\|_\infty \le C_1\left(\frac{t_0}{2}\right)^{-\frac{1}{2}}\|f\|_\infty, \qquad \forall\, t\ge t_0,$$

which is the desired second estimate. □

6. Application 3: modulus of continuity of transition semugroup of *log* α-stable processes

When the Lévy measure ν on $\mathbb{R}^d$ satisfies

$$\int_{\{|x|<r\}}\nu(\mathrm{d}x)=\infty, \qquad \forall\, r>0, \tag{6.1}$$

the process $(L_t)_{t\ge0}$ has infinite small jumps in any time interval $[t, t+\delta)$. Then (see e.g. [10]), the law $\mathfrak{L}(L_t)$ of L_t is continuous but not necessarily absolutely continuous with respect to Lebesgue measure. However, if additionally ν is absolutely continuous then $\mathfrak{L}(L_t)$ is absolutely continuous. Let us also recall, see e.g. [7], that the law $\mathfrak{L}(L_t)$ is absolutely continuous if and only if for the corresponding semigroup satisfies $P_t\colon B_b(\mathbb{R}^d)\mapsto C_b(\mathbb{R}^d)$. Thus the absolute continuity is equivalent to strong Feller property.

Below we provide some estimates for the moduli of continuity of the transition semigroup which is beyond the scope of α-stable type process. Namely, assume that the Lévy measure ν is absolutely continuous with respect to Lebesgue measure on the ball $B_1(0)=\{x\colon |x|<1\}$ and

$$\frac{\nu(\mathrm{d}x)}{\mathrm{d}x}\ge\frac{|\log_2|x||^{2\alpha}}{|x|^d}, \qquad x\in B_1(0), \tag{6.2}$$

where $\alpha\in(1,\infty)$ is a constant.

Theorem 6.1. *Let $(L_t)_{t\ge0}$ be a Lévy process with Lévy measure ν satisfying (6.2). Then $(L_t)_{t\ge0}$ is strong Feller. Moreover, there exists an $r_0>0$ such that*

$$|P_tf(x)-P_tf(y)|\le\frac{C}{|\log_2|x-y||^{\alpha-1}}, \qquad |x-y|\le r_0, \tag{6.3}$$

where C depends on α, d, t and r_0.

Let $\Omega\subset\mathbb{R}^d$ be an open set. Given a function $f\colon\Omega\to\mathbb{R}$, $x\in\Omega$ and $r>0$ such that the ball $B_r(x)\subset\Omega$, define

$$\bar f_{x,r}:=\frac{1}{|B_r(x)|}\int_{B_r(x)}f(y)\mathrm{d}y.$$

To show the theorem, we need to use the following lemma, which is a generalized Campanato theorem. The proof of the lemma is postponed to the appendix.

Lemma 6.2. *Let $\Omega\subset\mathbb{R}^d$ be open and bounded, and let $f\colon\Omega\to\mathbb{R}$ be a bounded function. Assume that there are constants $C>0$ and $\alpha>1$ such that*

$$\int_{B_r(x)}|f(y)-\bar f_{x,r}|^2\mathrm{d}y\le C\frac{r^d}{|\log_2 r|^{2\alpha}}, \tag{6.4}$$

for any ball $B_r(x) \subset \Omega$. Then f is uniformly continuous and there exists an $r_0 > 0$ such that for any open $\tilde{\Omega} \subset \Omega$ with $\operatorname{diam}(\tilde{\Omega}) < r_0$ and $\operatorname{dist}(\tilde{\Omega}, \partial\Omega) > r_0$, we have

$$\sup_{x,y\in\tilde{\Omega},x\neq y} |f(x) - f(y)||\log_2 |x-y||^{\alpha-1} \leq \tilde{C}, \tag{6.5}$$

where $\tilde{C}$ depends on C, $\operatorname{dist}(\tilde{\Omega}, \partial\Omega), d, \alpha$ and r_0.

Proof of Theorem 6.1. Let $f \in B_b(\mathbb{R}^d)$ and set $g(x) = P_t f(x)$. For any $t > 0$, we have

$$\begin{aligned}
\int_{B_r(x)} |g(y) - \bar{g}_{x,r}|^2 \mathrm{d}y &= \int_{\{|y-x|\leq r\}} \frac{1}{|B_r(x)|^2} \left| \int_{\{|z-x|\leq r\}} (g(y) - g(z))\, \mathrm{d}z \right|^2 \mathrm{d}y \\
&\leq \int_{\{|y-x|\leq r\}} \frac{1}{|B_r(x)|} \int_{\{|z-x|\leq r\}} |g(y) - g(z)|^2 \, \mathrm{d}z\mathrm{d}y \\
&\leq \int_{\{|y-x|\leq r\}} \frac{1}{|B_r(x)|} \int_{\{|y-z|\leq 2r\}} |g(y) - g(z)|^2 \, \mathrm{d}z\mathrm{d}y,
\end{aligned} \tag{6.6}$$

where the last inequality follows from the inclusion

$$\{|y-x| \leq r\} \cap \{|z-x| \leq r\} \subset \{|y-x| \leq r\} \cap \{|y-z| \leq 2r\}.$$

From (6.6) we further get

$$\begin{aligned}
&\int_{B_r(x)} |g(y) - \bar{g}_{x,r}|^2 \mathrm{d}y \\
&\leq \int_{\{|y-x|\leq r\}} \frac{1}{|B_r(x)|} \int_{\{|y-z|\leq 2r\}} |g(y) - g(z)|^2 \frac{|z-y|^d}{|\log_2 |y-z||^{2\alpha}} \frac{|\log_2 |z-y||^{2\alpha}}{|z-y|^d} \mathrm{d}z\mathrm{d}y \\
&= \int_{\{|y-x|\leq r\}} \frac{1}{|B_r(x)|} \int_{\{|z|\leq 2r\}} |g(y+z) - g(y)|^2 \frac{|z|^d}{|\log_2 |z||^{2\alpha}} \frac{|\log_2 |z||^{2\alpha}}{|z|^d} \mathrm{d}z\mathrm{d}y \\
&\leq \int_{\{|y-x|\leq r\}} \frac{1}{|B_r(x)|} \int_{\{|z|\leq 2r\}} |g(y+z) - g(y)|^2 \frac{|z|^d}{|\log_2 |z||^{2\alpha}} \nu(\mathrm{d}z)\mathrm{d}y
\end{aligned}$$

Since $\frac{r^d}{|\log_2 r|^{2\alpha}}$ is decreasing as $r < r_0/2$ for small $r_0 > 0$, the above inequality further gives

$$\begin{aligned}
\int_{B_r(x)} |g(y) - \bar{g}_{x,r}|^2 \mathrm{d}y &\leq C \frac{r^d}{|\log_2 r|^{2\alpha}} \frac{1}{|B_r(x)|} \int_{\{|y-x|\leq r\}} \int_{\{|z|\leq 2r\}} |g(y+z) - g(y)|^2 \, \nu(\mathrm{d}z)\mathrm{d}y \\
&\leq C \frac{r^d}{|\log_2 r|^{2\alpha}} \sup_{y\in B_r(x)} \int_{\mathbb{R}^d} |g(y+z) - g(y)|^2 \, \nu(\mathrm{d}z) \\
&\leq C \frac{r^d}{|\log_2 r|^{2\alpha}} t^{-1} \|f\|_\infty^2,
\end{aligned}$$

where the last inequality follows from Corollary 3.2. Combining the estimate above and Lemma 6.2 we obtain the desired conclusion. □

7. Appendix: Proof of Lemma 6.2

We shall follow [6]. For $0 < r_2 < r_1 < \min\{\text{dist}\,(\tilde{\Omega}, \partial\Omega), 1\}$ and any $x \in \tilde{\Omega}$, we have

(7.1)
$$\begin{aligned}
|\bar{f}_{x,r_1} - \bar{f}_{x,r_2}| &\le \frac{1}{|B_{r_2}|}\int_{B_{r_2}(x)} |f(y) - \bar{f}_{x,r_2}|\mathrm{d}y + \frac{1}{|B_{r_2}|}\int_{B_{r_2}(x)} |f(y) - \bar{f}_{x,r_1}|\mathrm{d}y \\
&\le \sqrt{\frac{1}{|B_{r_2}|}\int_{B_{r_2}(x)} |f(y) - \bar{f}_{x,r_2}|^2\mathrm{d}y} + \sqrt{\frac{1}{|B_{r_2}|}\int_{B_{r_2}(x)} |f(y) - \bar{f}_{x,r_1}|^2\mathrm{d}y} \\
&\le \sqrt{\frac{1}{|B_{r_2}|}\int_{B_{r_2}(x)} |f(y) - \bar{f}_{x,r_2}|^2\mathrm{d}y} + \sqrt{\frac{|B_{r_1}|}{|B_{r_2}|}\frac{1}{|B_{r_1}|}\int_{B_{r_1}(x)} |f(y) - \bar{f}_{x,r_1}|^2\mathrm{d}y} \\
&\le C\left[\frac{1}{|\log_2 r_2|^\alpha} + \left(\frac{r_1}{r_2}\right)^{d/2}\frac{1}{|\log_2 r_1|^\alpha}\right] \\
&\le C\left[1 + \left(\frac{r_1}{r_2}\right)^{d/2}\right]\frac{1}{|\log_2 r_1|^\alpha},
\end{aligned}$$

where the last inequality is by the assumption of the lemma. For all $0 < r_n < r_m < \text{dist}\,(\tilde{\Omega}, \partial\Omega)$, define

$$N := \left[\log_2\left(\frac{r_m}{r_n}\right)\right],$$

without loss of generality we assume $r_m < 1/2$. By (7.1) we have

(7.2)
$$\begin{aligned}
|\bar{f}_{x,r_n} - \bar{f}_{x,r_m}| &\le \sum_{k=1}^{N} |\bar{f}_{x,2^{-k}r_m} - \bar{f}_{x,2^{-k+1}r_m}| + |\bar{f}_{x,2^{-N}r_m} - \bar{f}_{x,r_n}| \\
&\le C_d \sum_{k=1}^{N} \frac{1}{|k - 1 - \log_2 r_m|^\alpha} + C_d \frac{1}{|N - \log_2 r_m|^\alpha} \\
&\le C_{d,\alpha} \frac{1}{|\log_2 r_m|^{\alpha-1}}.
\end{aligned}$$

Hence, there exists an $\tilde{f}$ such that

$$\lim_{r\to 0} \bar{f}_{x,r} = \tilde{f}(x), \qquad \forall\, x \in \tilde{\Omega},$$

and there exists an $r_0 > 0$ such that as $r < r_0$,

$$|\bar{f}_{x,r} - \tilde{f}(x)| \le C_{d,\alpha} \frac{1}{|\log_2 r|^{\alpha-1}}, \qquad \forall\, x \in \tilde{\Omega}. \tag{7.3}$$

On the other hand, by the Lebesgue theorem,

$$\lim_{r\to 0} \bar{f}_{x,r} = f(x), \qquad \text{a.s. } x \in \tilde{\Omega}.$$

By (7.3), all the points in $\tilde{\Omega}$ are Lebesgue points. Hence, $\bar{f}_{x,r} \to f(x)$ uniformly for $x \in \tilde{\Omega}$ as $r \to 0$ with

$$|\bar{f}_{x,r} - f(x)| \le C_{d,\alpha} \frac{1}{|\log_2 r|^{\alpha-1}}, \qquad \forall\, x \in \tilde{\Omega}. \tag{7.4}$$

Now for $x, y \in \tilde{\Omega}$, denote $r = |x - y|$, we have

$$|\bar{f}_{y,2r} - \bar{f}_{x,2r}| \le |\bar{f}_{y,2r} - f(z)| + |f(z) - \bar{f}_{x,2r}|, \qquad z \in B_{2r}(x) \cap B_{2r}(y). \tag{7.5}$$

Since $B_{2r}(x) \cap B_{2r}(y)$ contains a ball with radius r, as $r < r_0/2$,

$$
\begin{aligned}
|\bar{f}_{y,2r} - \bar{f}_{x,2r}| &\le \frac{1}{|B_r|} \int_{B_{2r}(x)\cap B_{2r}(y)} |\bar{f}_{y,2r} - f(z)| + |f(z) - \bar{f}_{x,2r}| \mathrm{d}z \\
&\le \frac{1}{|B_r|} \int_{B_{2r}(y)} |\bar{f}_{y,2r} - f(z)| \mathrm{d}z + \frac{1}{|B_r|} \int_{B_{2r}(x)} |f(z) - \bar{f}_{x,2r}| \mathrm{d}z \\
&\le C \frac{1}{|\log_2 r|^\alpha},
\end{aligned} \tag{7.6}
$$

where the last inequality is by the assumption of the lemma. Observe that

$$
|f(x) - f(y)| \le |f(x) - \bar{f}_{x,2r}| + |f(y) - \bar{f}_{y,2r}| + |\bar{f}_{y,2r} - \bar{f}_{x,2r}|. \tag{7.7}
$$

This, together with (7.6) and (7.4), immediately implies the desired inequality.

References

1. D. Applebaum, *Lévy Processes and Stochastic Calculus*, Cambridge Studies in Advance Mathematics 93, Cambridge University Press, 2004.
2. J. Bao, F.Y. Wang and C. Yuan, *Bismut formulae and applications for functional SPDEs*, Bull. Sci. Math. 137 (2013), no. 4, 509–522.
3. Z.Q. Chen and J. Wang, *Perturbation by non-local operators*, preprint.
4. K.D. Elworthy and X.M. Li, *Formulae for the derivatives of heat semigroups*, J. Funct. Anal. 125, no. 1, (1994), 252–286.
5. J.P. Eckmann and M. Hairer, *Uniqueness of the invariant measure for a stochastic PDE driven by degenerate noise*, Commun. Math. Phys. 219 (2001), no 3, pp. 523–565
6. Q. Han and F. Lin, *Elliptic Partial Differential Equations*, second ed., Courant Lecture Notes in Mathematics, Vol. 1, Courant Institute of Mathematical Sciences, New York, 2011. MR 2777537 (2012c:35077)
7. J. Hawkes, *Potential theory of Léve processes*, Proc. London Math. Soc. 38 (1979) 335–352.
8. S. Peszat and J. Zabczyk, *Strong Feller property and irreducibility for diffusions on Hilbert spaces*, Ann. Probab. 23 (1995), 157–172.
9. S. Peszat and J. Zabczyk, *Stochastic Partial Differential Equations with Lévy Noise (an Evolution Equation Approach)*, Cambridge University Press, Cambridge, 2007.
10. K. Sato, *Lévy Processes and Infinitely Divisible Distributions*, Cambridge Studies in Advanced Mathematics, 68. Cambridge University Press, Cambridge, 1999. xii+486 pp.
11. A. Takeuchi, *Bismut-Elworthy-Li type formulae for stochastic differential equations with jumps*, J. Theoretical Probab., 23 (2010), no. 2, 576–604.
12. F.Y. Wang, *Gradient estimate for Ornstein-Uhlenbeck jump processes*, Stoch. Proc. Appl. 121 (2011), no. 3, 466–478.
13. F.Y. Wang, *Integration by parts formula and shift Harnack inequality for stochastic equations*, Ann. Probab. 42 (2014), no. 3, 994–1019.
14. F.Y. Wang and L. Xu, *Derivative formula and applications for hyperdissipative stochastic Navier-Stokes/Burgers equations*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. 15 (2012), no. 3, 1250020, 19 pp.
15. F.Y. Wang, L. Xu and X. Zhang: *Gradient estimates for SDEs driven by multiplicative Levy noise*, arXiv 1301.4528.
16. F.Y. Wang and L. Yan, *Gradient estimate on convex domains and applications*, Proc. Amer. Math. Soc. 141 (2013), no. 3, 1067–1081
17. X. Zhang, *Derivative formulas and gradient estimates for SDEs driven by α-stable processes*, Stochastic Process. Appl. 123 (2013), no. 4, 1213–1228.

Institute of Applied Mathematics, Academy of Mathematics and Systems Sciences, Academia Sinica, P.R.China
E-mail address: dzhao@amt.ac.cn

Institute of Mathematics, Polish Academy of Sciences, Św. Tomasza 30/7, 31-027 Cracow, Poland
E-mail address: napeszat@cyf-kr.edu.pl

Department of Mathematics, Faculty of Science and Technology University of Macau, Av. Padre Tomás Pereira, Taipa Macau, China
E-mail address: lihuxu@umac.mo